



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Middle Tropospheric CO₂ and O₃ by the Atmospheric Infrared Sounder

Xun Jiang, Moustafa T. Chahine, Edward T. Olsen, Qinbin Li,
Luke Chen, Thomas Pagano, and Yuk Yung

Jet Propulsion Laboratory
California Institute of Technology

04/15/2008



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Overview

- **Motivation**
- **Validation of AIRS CO₂**
- **Large-scale dynamics at the polar region**
- **Semi-Annual Oscillation in CO₂**



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Motivation

- Polar regions have profound significance for climate, however, there are a lot of unknown due to limited observations.
- Current models cannot work well in the polar region.
- One important issue is to simulate the exchange between stratosphere and troposphere correctly.



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Data

➤ AIRS CO₂ and O₃

CO₂ Weighting Function



O₃ Weighting Function



CO₂ maximum sensitivity is at 500-300 hPa.

O₃ maximum sensitivity is at 300 hPa.



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder

Data

- AIRS CO₂ and O₃
- Aircraft Data of CO₂ from *Matsueda et al.* [2002], Climate Monitoring & Diagnostics Laboratory (CMDL), and SPURT Aircraft [*Hoor et al.*, 2004]
- Ozonesonde data from World Ozone and Ultraviolet Data (WOUDC)


Version 5 VPD AIRS CO₂

01–30 Apr 2003



Version 5 VPD AIRS O₃


01–30 Apr 2003



Stratospheric Sudden Warming (SSW)

- Strongest dynamical coupling between stratosphere and troposphere
- During SSW:
Polar stratospheric temperature rise and the circum-polar flow reverse direction in a few days
- After SSW:
Decrease of vortex area; Less downwelling in the polar region
- Important influence on chemical tracers

Influence of Sudden Warming on Tracers




After SSW:

***Decrease of Vortex Area -----> High CO₂ is transported into Pole
Low O₃ is transported into Pole***

Polar CO₂ should increase and polar O₃ should decrease after the final warming

Influence of Sudden Warming on Tracers




After SSW:

***Less Downwelling -----> Less stratospheric low CO₂ (high O₃)
will be transported into troposphere***


Mid trop CO₂ should increase and mid trop O₃ should decrease after the final warming

Influence of Sudden Stratospheric Warming on CO₂ and O₃ AIRS- April 2003



AIRS retrieved upper tropospheric CO₂ increases while AIRS 300 mb O₃ decreases following a sudden stratospheric warming event

Influence of Sudden Stratospheric Warming on CO₂ and O₃ AIRS- April 2003




AIRS retrieved upper tropospheric CO₂ increases while AIRS 300 mb O₃ decreases following a sudden stratospheric warming event

Before SSW


(a) AIRS CO₂ in Apr 1–10

CO₂




After SSW

(b) AIRS CO₂ in Apr 21–30




(c) AIRS O₃ in Apr 1–10

O₃




(d) AIRS O₃ in Apr 21–30



Contour: NCEP2 GPH at 500 hPa


AIRS CO₂ (Apr 2003)

AIRS NP Mid-Tropospheric CO₂. April 2003, V5 Day 3 x 5



AIRS O₃ (Apr 2003)


AIRS NP 300mb O₃. April 2003, V5 Day 3 x 5



Overview


- Motivation
- Validation of AIRS CO₂
- Stratospheric Sudden Warming: Influence on CO₂ and O₃
- **Semi-Annual Oscillation in CO₂**

Semi-Annual Oscillation in Aircraft and Model CO₂





● Matsueda CO₂ aircraft data — GEOS-Chem 3D; G-4 (CMDL BC) — GEOS-Chem 3D; G-4 (Source/Sink)
 — CJCTM 2D (CMDL BC) — MOZART2 (CMDL BC)

Semi-Annual Oscillation in Aircraft and Model CO₂



● Matsueda CO₂ aircraft data — GEOS-Chem 3D; G-4 (CMDL BC) — GEOS-Chem 3D; G-4 (Source/Sink)
 — CJCTM 2D (CMDL BC) — MOZART2 (CMDL BC)


Semi-Annual Oscillation in AIRS CO₂




- Matsueda CO₂ aircraft data — GEOS-Chem 3D; G-4 (CMDL BC) — GEOS-Chem 3D; G-4 (Source/Sink)
- CJCTM 2D (CMDL BC) — MOZART2 (CMDL BC)
- Version 5 AIRS CO₂

Source for Semi-Annual Oscillation


NPP = Net Primary Production (Photosynthesis)



RH = Respiration



NEP = RH + NPP



**Sum of the net primary production and respiration from biosphere
lead to the Semi-Annual Oscillation in CO_2**

Conclusions

- With AIRS, we monitor the distribution and transport of global CO₂ on a weekly basis for the first time.
- The latitudinal distribution of AIRS retrievals of upper tropospheric CO₂ agrees reasonably well with in situ aircraft observations of CO₂ and model simulations.
- AIRS retrieved upper tropospheric CO₂ increases while AIRS 300 mb O₃ decreases following a sudden stratospheric warming event.
- Semi-annual Oscillation is found in the upper tropospheric CO₂.



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Thank you!